arXiv:1009.1718v1 [math.DG] 9 Sep 2010

SOME SUBMANIFOLDS OF ALMOST CONTACT MANIFOLDS WITH NORDEN METRIC *

G.NAKOVA

Department of Algebra and Geometry,
Faculty of Pedagogics, University of Veliko Tarnovo,
1 Theodosij Tirnovsky Str., 5000 Veliko Tarnovo, Bulgaria
E-mail: gnakova@yahoo.com

In this paper we study submanifolds of almost contact manifolds with Norden metric of codimension two with totally real normal spaces. Examples of such submanifolds as a Lie subgroups are constructed.

Keywords: Norden metric, almost contact manifold, submanifold, Lie group

1. Introduction

Let $(M, \varphi, \xi, \eta, g)$ be a $(2n+1)$-dimensional almost contact manifold with Norden metric, i. e. (φ, ξ, η) is an almost contact structure[1] and g is a metric[3] on M such that

$$\begin{aligned} &\varphi^2 X = -id + \eta \otimes \xi, \qquad \eta(\xi) = 1, \\ &g(\varphi X, \varphi Y) = -g(X,Y) + \eta(X)\eta(Y), \end{aligned} \tag{1}$$

where id denotes the identity transformation and X, Y are differentiable vector fields on M, i. e. $X, Y \in \mathfrak{X}(M)$. The tensor $\tilde{g}$ given by $\tilde{g}(X,Y) = g(X, \varphi Y) + \eta(X)\eta(Y)$ is a Norden metric, too. Both metrics g and $\tilde{g}$ are indefinite of signature $(n+1, n)$.

Let ∇ be the Levi-Civita connection of the metric g. The tensor field F of type $(0,3)$ on M is defined by

$$F(X,Y,Z) = g((\nabla_X \varphi)Y, Z).$$

A classification of the almost contact manifolds with Norden metric with respect to the tensor F is given in[3] and eleven basic classes $\mathcal{F}_i (i = 1, 2, \ldots, 11)$ are obtained.

---

*Partially supported by Scientific researches fund of "St. Cyril and St. Methodius" University of Veliko Tarnovo under contract RD-491-08 from 27. 06. 2008.

Let R be the curvature tensor field of ∇ defined by

$$R(X,Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z.$$

The corresponding tensor field of type $(0,4)$ is determined as follows

$$R(X,Y,Z,W) = g(R(X,Y)Z,W).$$

Let $(\overline{M}, \overline{\varphi}, \overline{\xi}, \overline{\eta}, g)$ $(dim\overline{M} = 2n+3)$ be an almost contact manifold with Norden metric and let M be a submanifold of $\overline{M}$. Then for each point $p \in \overline{M}$ we have

$$T_p\overline{M} = T_pM \oplus (T_pM)^{\perp},$$

where T_pM and $(T_pM)^{\perp}$ are the tangent space and the normal space of $\overline{M}$ at p respectively. When the submanifold M of $\overline{M}$ is of codimension 2 we denote $(T_pM)^{\perp}$ by $\alpha = \{N_1, N_2\}$, i.e. α is a normal section of M.

Let α be a 2-dimensional section in $T_p\overline{M}$. Let us recall a section α is said to be

- *non-degenerate*, *weakly isotropic* or *strongly isotropic* if the rank of the restriction of the metric g on α is 2, 1 or 0 respectively;
- of *pure* or *hybrid* type if the restriction of g on α has a signature $(2,0)$, $(0,2)$ or $(1,1)$ respectively;
- *holomorphic* if $\overline{\varphi}\alpha = \alpha$;
- $\overline{\xi}$- section if $\overline{\xi} \in \alpha$;
- *totally real* if $\overline{\varphi}\alpha \perp \alpha$.

Submanifolds M of $\overline{M}$ of codimension 2 with a non-degenerate of hybrid type normal section α are studied. In[4] two basic types of such submanifolds are considered: α is a holomorphic section and α is a $\overline{\xi}$- section. In[5] the normal section $\alpha = \{N_1, N_2\}$ is such that $\overline{\varphi}N_1 \notin \alpha$, $\overline{\varphi}N_2 \in \alpha$. In this paper we consider submanifolds M of $\overline{M}$ of codimension 2 in the case when the normal section α is a non-degenerate of hybrid type and α is a totally real. The totally real sections α are two types: α is non-orthogonal to $\overline{\xi}$ and α is orthogonal to $\overline{\xi}$.

2. Submanifolds of codimension 2 of almost contact manifolds with Norden metric with totally real non-orthogonal to $\overline{\xi}$ normal spaces

Let $(\overline{M}, \overline{\varphi}, \overline{\xi}, \overline{\eta}, g)$ $(dim\overline{M} = 2n+3)$ be an almost contact manifold with Norden metric and let M be a submanifold of codimension 2 of $\overline{M}$. We assume that there exists a normal section $\alpha = \{N_1, N_2\}$ defined globally over the submanifold M such that

- α is a non-degenerate of hybrid type, i.e.

$$g(N_1,N_1)=-g(N_2,N_2)=1,\quad g(N_1,N_2)=0; \tag{2}$$

- α is a totally real, i.e.

$$g(N_1,\overline{\varphi}N_1)=g(N_2,\overline{\varphi}N_2)=g(N_1,\overline{\varphi}N_2)=g(N_2,\overline{\varphi}N_1)=0; \tag{3}$$

- α is a non-orthogonal to $\overline{\xi}$ ($\overline{\xi}\notin T_pM$) and $\overline{\xi}\notin\alpha$.

Then we obtain the following decomposition for $\overline{\xi}$, $\overline{\varphi}X$, $\overline{\varphi}N_1$, $\overline{\varphi}N_2$ with respect to $\{N_1,N_2\}$ and T_pM

$$\begin{array}{l} \overline{\xi}=\xi_0+aN_1+bN_2;\\ \overline{\varphi}X=\varphi X+\eta^1(X)N_1+\eta^2(X)N_2,\ X\in\chi(M);\\ \overline{\varphi}N_1=\xi_1;\\ \overline{\varphi}N_2=-\xi_2; \end{array} \tag{4}$$

where φ denotes a tensor field of type $(1,1)$ on M; $\xi_0,\xi_1,\xi_2\in\chi(M)$; η^1 and η^2 are 1-forms on M; a,b are functions on M such that $(a,b)\neq(0,0)$. We denote the restriction of g on M by the same letter.

Let $a\neq 0$, $|a|>b$ and $a^2-b^2=k^2$. Taking into account the equalities $(1)\div(4)$ we compute

$$\eta^i(X)=g(X,\xi_i),\ (i=0,1,2); \tag{5}$$

$$g(\varphi X,\varphi Y)=-g(X,Y)+\eta^0(X)\eta^0(Y)-\eta^1(X)\eta^1(Y)+\eta^2(X)\eta^2(Y); \tag{6}$$

$$\begin{array}{l} \varphi^2X=-X+\eta^0(X)\xi_0-\eta^1(X)\xi_1+\eta^2(X)\xi_2;\\ \eta^0(\varphi X)=-a\eta^1(X)+b\eta^2(X);\\ \eta^1(\varphi X)=a\eta^0(X);\quad \eta^2(\varphi X)=b\eta^0(X); \end{array} \tag{7}$$

$$\varphi\xi_0=-a\xi_1+b\xi_2;\quad \varphi\xi_1=a\xi_0;\quad \varphi\xi_2=b\xi_0; \tag{8}$$

$$\begin{array}{l} g(\xi_0,\xi_0)=1-a^2+b^2;\quad g(\xi_1,\xi_1)=a^2-1;\\ g(\xi_2,\xi_2)=1+b^2;\quad g(\xi_0,\xi_1)=g(\xi_0,\xi_2)=0;\\ g(\xi_1,\xi_2)=ab; \end{array} \tag{9}$$

for arbitrary $X,Y\in\chi(M)$.

Now we define a vector field ξ, an 1-form η and a tensor field ϕ of type $(1,1)$ on M by

$$\begin{aligned} &\xi = -\frac{b}{k}.\xi_1 + \frac{a}{k}.\xi_2; \\ &\eta(X) = -\frac{b}{k}.\eta^1(X) + \frac{a}{k}.\eta^2(X), \quad X \in \chi(M); \\ &\phi X = \lambda.\varphi^3 X + \mu.\varphi X, \quad X \in \chi(M); \end{aligned} \tag{10}$$

where

$$\lambda_1 = \frac{\epsilon}{k(k+1)}, \quad \mu_1 = \frac{\epsilon(1+k^2+k)}{k(k+1)};$$

$$\lambda_2 = \frac{\epsilon}{k(k-1)}, \quad \mu_2 = \frac{\epsilon(1+k^2-k)}{k(k-1)}; \quad \epsilon = \pm 1.$$

Further we consider the following cases for k:

1) $k^2 \neq 1 \iff k \neq \pm 1$. In this case ϕ, ξ, η are given by (10) and $\lambda = \lambda_1$, $\mu = \mu_1$ or $\lambda = \lambda_2$, $\mu = \mu_2$.

2) $k = -1$. We obtain ϕ, ξ, η from (10) by $k = -1$ and $\lambda = \lambda_2 = \dfrac{\epsilon}{2}$, $\mu = \mu_2 = \dfrac{3\epsilon}{2}$.

3) $k = 1$. We obtain ϕ, ξ, η from (10) by $k = 1$ and $\lambda = \lambda_1 = \dfrac{\epsilon}{2}$, $\mu = \mu_1 = \dfrac{3\epsilon}{2}$.

Using $(5) \div (10)$ we verify that (ϕ, ξ, η) is an almost contact structure on M and the restriction of g on M is Norden metric. Thus, the submanifolds (M, ϕ, ξ, η, g) of $\overline{M}$ considered in 1), 2), 3) are $(2n+1)$-dimensional almost contact manifolds with Norden metric.

Denoting by $\overline{\nabla}$ and ∇ the Levi-Civita connections of the metric g in $\overline{M}$ and M respectively, the formulas of Gauss and Weingarten are

$$\begin{aligned} &\overline{\nabla}_X Y = \nabla_X Y + g(A_{N_1}X, Y)N_1 - g(A_{N_2}X, Y)N_2; \\ &\overline{\nabla}_X N_1 = -A_{N_1}X + \gamma(X)N_2; \\ &\overline{\nabla}_X N_2 = -A_{N_2}X + \gamma(X)N_1, \quad X, Y \in \chi(M); \end{aligned} \tag{11}$$

where A_{N_i} $(i = 1, 2)$ are the second fundamental tensors and γ is an 1-form on M.

3. Submanifolds of codimension 2 of almost contact manifolds with Norden metric with totally real orthogonal to $\overline{\xi}$ normal spaces

Let $(\overline{M}, \overline{\varphi}, \overline{\xi}, \overline{\eta}, g)$ $(dim\overline{M} = 2n+3)$ be an almost contact manifold with Norden metric and let M be a submanifold of codimension 2 of $\overline{M}$. We assume that there exists a normal section $\alpha = \{N_1, N_2\}$ defined globally over the submanifold M such that

- α is a non-degenerate of hybrid type, i.e. the equality (2) holds;
- α is a totally real;
- α is orthogonal to $\overline{\xi}$, i.e. $\overline{\xi} \in T_pM$.

Then from (4) by $a = b = 0$ we obtain the following decomposition with respect to $\{N_1, N_2\}$ and T_pM

$$\begin{aligned}
&\overline{\xi} = \xi_0;\\
&\overline{\varphi}X = \varphi X + \eta^1(X)N_1 + \eta^2(X)N_2,\ X \in \chi(M);\\
&\overline{\varphi}N_1 = \xi_1;\\
&\overline{\varphi}N_2 = -\xi_2.
\end{aligned} \tag{12}$$

Substituting $a = b = 0$ in $(7), (8), (9)$ we have

$$\begin{aligned}
&\eta^0(\varphi X) = \eta^1(\varphi X) = \eta^2(\varphi X) = 0;\\
&\varphi\xi_0 = \varphi\xi_1 = \varphi\xi_2 = 0;\\
&g(\xi_0, \xi_0) = g(\xi_2, \xi_2) = 1;\ g(\xi_1, \xi_1) = -1;\\
&g(\xi_0, \xi_1) = g(\xi_0, \xi_2) = g(\xi_1, \xi_2) = 0.
\end{aligned} \tag{13}$$

Now we define a vector field ξ, an 1-form η and a tensor field ϕ of type $(1,1)$ on M by

$$\begin{aligned}
&\xi = t_0\xi_0 - t_2\xi_2;\\
&\eta(X) = t_0\eta^0(X) - t_2\eta^2(X), \quad X \in \chi(M);\\
&\phi X = \varphi X + t_0\{\eta^1(X).\xi_2 + \eta^2(X).\xi_1\} +\\
&t_2\{\eta^0(X).\xi_1 + \eta^1(X).\xi_0\};
\end{aligned} \tag{14}$$

where t_0, t_2 are functions on M and $t_0^2 + t_2^2 = 1$.

Using $(5), (6), (13), (14)$ we verify that (ϕ, ξ, η) is an almost contact structure on M and the restriction of g on M is Norden metric. So, the submanifolds (M, ϕ, ξ, η, g) of $\overline{M}$ are $(2n+1)$-dimensional almost contact manifolds with Norden metric. The formulas of Gauss and Weingarten are the same as those in section 2.

4. Examples of submanifolds of codimension 2 of almost contact manifolds with Norden metric with totally real normal spaces

In[6] a Lie group as a 5-dimensional almost contact manifold with Norden metric of the class $\mathcal{F}_9$ is constructed. We will use this Lie group to obtain examples of submanifolds considered in sections 2 and 3.

First we recall some facts from[6] which we need. Let $\mathfrak{g}$ be a real Lie algebra with a global basis of left invariant vector fields $\{X_1, X_2, X_3, X_4, X_5\}$ and G be the associated with $\mathfrak{g}$ real connected Lie group. The almost contact structure $(\overline{\varphi}, \overline{\xi}, \overline{\eta})$ and the Norden metric g on G are defined by:

$$\begin{array}{l}
\varphi X_i = X_{2+i}, \ \ \varphi X_{2+i} = -X_i, \ \ \varphi X_5 = 0, \ \ (i = 1, 2); \\
g(X_i, X_i) = -g(X_{2+i}, X_{2+i}) = g(X_5, X_5) = 1, \ \ (i = 1, 2); \\
g(X_j, X_k) = 0, \quad (j \neq k, \ \ j, k = 1, 2, 3, 4, 5); \\
\overline{\xi} = X_5, \quad \overline{\eta}(X_i) = g(X_i, X_5), \ (i = 1, 2, 3, 4, 5).
\end{array} \tag{15}$$

The commutators of the basis vector fields are given by:

$$\begin{array}{l}
[X_1, X_2] = -[X_1, X_3] = aX_4, \quad [X_2, X_3] = aX_2 + aX_3, \\
\\
[X_3, X_4] = -[X_2, X_4] = aX_1, \quad [X_2, \overline{\xi}] = 2mX_1, \\
\\
[X_3, \overline{\xi}] = -2mX_4, \quad [X_1, X_4] = [X_1, \overline{\xi}] = [X_4, \overline{\xi}] = 0, \\
a, m \in \mathcal{R}.
\end{array} \tag{16}$$

So, the manifold $(G, \overline{\varphi}, \overline{\xi}, \overline{\eta}, g)$ is an almost contact manifold with Norden metric in the class $\mathcal{F}_9$.

Theorem 4.1. [2] *Let G be a Lie group with a Lie algebra $\mathfrak{g}$ and $\widetilde{b}$ be a subalgebra of $\mathfrak{g}$. There exists an unique connected Lie subgroup H of G such that the Lie algebra b of H coincides with $\widetilde{b}$.*

From the equalities for the commutators of the basis vector fields $\{X_1, X_2, X_3, X_4, X_5\}$ it follows that the 3-dimensional subspaces of $\mathfrak{g}$ $\mathbf{b_1}$ with a basis $\{X_1, X_2, X_3\}$, $\mathbf{b_2}$ with a basis $\{X_1, X_3, X_4\}$ and $\mathbf{b_3}$ with a basis $\{X_1, X_4, \overline{\xi}\,\}$ are closed under the bracket operation. Hence $b_i \ \ (i = 1, 2, 3)$ are real subalgebras of $\mathfrak{g}$. Taking into account Theorem 4.1 we have there exist Lie subgroups H_i $(i = 1, 2, 3)$ of the Lie group G with Lie algebras b_i $(i = 1, 2, 3)$ respectively. The normal spaces α_i $(i = 1, 2, 3)$ of the submanifolds H_i $(i = 1, 2, 3)$ of G are: $\alpha_1 = \{X_4, \overline{\xi}\}$, $\alpha_2 = \{X_2, \overline{\xi}\}$, $\alpha_3 = \{X_2, X_3\}$. Because of (15) we have α_1 is $\overline{\xi}$- section of hybrid type, α_2 is $\overline{\xi}$- section of pure type and α_3 is a totally real orthogonal to $\overline{\xi}$ section of hybrid type.

So, the submanifold H_3 of G is of the same type submanifolds considered in section 3.

We choose the unit normal fields of H_3 $N_1 = X_2$ and $N_2 = X_3$. For an arbitrary $X \in \chi(H_3)$ we have $X = x^1 X_1 + x^4 X_4 + \overline{\eta}(X)\overline{\xi}$. Taking into account (15) we compute

$$\begin{aligned} &\overline{\xi} = \xi_0; \\ &\overline{\varphi} X = -x^4 X_2 + x^1 X_3; \\ &\overline{\varphi} X_2 = X_4; \\ &\overline{\varphi} X_3 = -X_1. \end{aligned} \tag{17}$$

From (12), (17) it follows

$$\begin{aligned} &\eta^0(X) = \overline{\eta}(X); \\ &\varphi X = 0;\ \eta^1(X) = -x^4;\ \eta^2(X) = x^1; \\ &\xi_1 = X_4;\ \xi_2 = X_1. \end{aligned} \tag{18}$$

Substituting (18) in (14) for the almost contact structure on H_3 we obtain

$$\begin{aligned} &\xi = t_0 \overline{\xi} - t_2 X_1; \\ &\eta(X) = t_0 \overline{\eta}(X) - t_2 x^1; \\ &\phi X = t_0\{-x^4 X_1 + x^1 X_4\} + t_2\{\overline{\eta}(X) X_4 - x^4 \overline{\xi}\}; \end{aligned} \tag{19}$$

where $t_0,\ t_2 \in \mathcal{R}$ and $t_0^2 + t_2^2 = 1$.

Using the well known condition for the Levi-Civita connection ∇ of g

$$\begin{aligned} 2g(\nabla_X Y, Z) = {} & Xg(Y,Z) + Yg(X,Z) - Zg(X,Y) + g([X,Y],Z) \\ & + g([Z,X],Y) + g([Z,Y],X) \end{aligned} \tag{20}$$

we get the following equation for the tensor F of H_3

$$\begin{aligned} F(X,Y,Z) = {} & \frac{1}{2}\{g([X,\phi Y] - \phi[X,Y], Z) \\ & +g(\phi[Z,X] - [\phi Z, X], Y) \\ & +g([Z,\phi Y] - [\phi Z, Y], X)\}, \quad X,Y,Z \in \chi(H_3). \end{aligned} \tag{21}$$

From (16) we have $[X_1, X_4] = [X_1, \overline{\xi}] = [X_4, \overline{\xi}] = 0$. Having in mind the last equalities, (19) and (21) for the tensor F of H_3 we obtain $F = 0$. Thus, the submanifold $(H_3, \phi, \xi, \eta, g)$ of G, where (ϕ, ξ, η) is defined by (19) is an almost contact manifold with Norden metric in the class $\mathcal{F}_0$.

In order to construct an example for a submanifold from section 2 we make the following change of the basis of $\mathfrak{g}$

$$\begin{pmatrix} E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \end{pmatrix} = T^T \begin{pmatrix} X_1 \\ X_2 \\ \overline{\xi} \\ X_3 \\ X_4 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{\sqrt{3}}{2} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \in O(3,2). \tag{22}$$

Taking into account (16) and (22) we compute the commutators of the basis vector fields $\{E_1, E_2, E_3, E_4, E_5\}$ of $\mathfrak{g}$

$$\begin{aligned} &[E_1, E_2] = \frac{\sqrt{3}}{2} a E_5, \quad [E_1, E_3] = \frac{1}{2} a E_5, \quad [E_3, E_5] = -\frac{1}{2} a E_1, \\ &[E_2, E_5] = -\frac{\sqrt{3}}{2} a E_1, \quad [E_1, E_4] = -a E_5, \\ &[E_2, E_4] = \frac{3}{4} a E_2 + \frac{\sqrt{3}}{4} a E_3 + \frac{\sqrt{3}}{2} a E_4 - m E_5, \\ &[E_3, E_4] = \frac{\sqrt{3}}{4} a E_2 + \frac{1}{4} a E_3 + \frac{1}{2} a E_4 + \sqrt{3} m E_5, \\ &[E_4, E_5] = a E_1, \quad [E_2, E_3] = 2m E_1, \quad [E_1, E_5] = 0. \end{aligned} \tag{23}$$

Because of the elements of the matrix T are constants the Jacobi identity for the vector fields $\{E_1, E_2, E_3, E_4, E_5\}$ is valid. Now, we compute the matrix B of $\overline{\varphi}$ and the coordinates of $\overline{\xi}$ with respect to the basis $\{E_1, E_2, E_3, E_4, E_5\}$

$$B = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -\frac{\sqrt{3}}{2} \\ 0 & 0 & 0 & 0 & -\frac{1}{2} \\ 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix}; \quad \overline{\xi} = \left(0, -\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, 0\right). \tag{24}$$

From $T \in O(3,2)$ it follows the matrix of the metric g with respect to the basis $\{E_1, E_2, E_3, E_4, E_5\}$ is the same as the matrix

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix} \tag{25}$$

of g with respect to the basis $\{X_1, X_2, \overline{\xi}, X_3, X_4\}$.

Using (23) we have that the 3-dimensional subspace b of $\mathfrak{g}$ with a basis $\{E_1, E_2, E_5\}$ is a subalgebra of $\mathfrak{g}$. Let H be the Lie subgroup of G with a Lie algebra b. Having in mind (24), (25) we obtain that the section $\alpha = \{E_3, E_4\}$ is a normal to the submanifold H, α is a totally real non-orthogonal to $\overline{\xi}$ section of hybrid type and $\overline{\xi} \notin \alpha$, i.e. H is of the same type submanifolds considered in section 2. We have the following decomposition of $\overline{\xi}, \overline{\varphi}X, \overline{\varphi}E_3, \overline{\varphi}E_4$ with respect to $\{E_3, E_4\}$ and T_pH

$$\begin{aligned}
&\overline{\xi} = -\frac{1}{2}E_2 + \frac{\sqrt{3}}{2}E_3;\\
&\overline{\varphi}X = -\frac{\sqrt{3}}{2}\overline{x}^5 E_2 + \frac{\sqrt{3}}{2}\overline{x}^2 E_5 - \frac{1}{2}\overline{x}^5 E_3 + \overline{x}^1 E_4;\\
&\overline{\varphi}E_3 = \frac{1}{2}E_5;\\
&\overline{\varphi}E_4 = -E_1;
\end{aligned} \tag{26}$$

where $X \in \chi(H)$ and $X = \overline{x}^1 E_1 + \overline{x}^2 E_2 + \overline{x}^5 E_5$. We substitute

$$a = \frac{\sqrt{3}}{2}, \quad b = 0, \quad k = \frac{\sqrt{3}}{2}, \quad \lambda = \frac{4\sqrt{3}(2-\sqrt{3})}{3}, \quad \mu = \lambda + 1,$$

$$\xi_2 = E_1, \quad \eta^2(X) = \overline{x}^1, \quad \varphi X = -\frac{\sqrt{3}}{2}\overline{x}^5 E_2 + \frac{\sqrt{3}}{2}\overline{x}^2 E_5$$

in (10) and obtain an almost contact structure (ϕ, ξ, η)

$$\begin{aligned}
&\xi = E_1;\\
&\eta^(X) = \overline{x}^1;\\
&\phi X = \frac{2\sqrt{3}}{3}\varphi X = -\overline{x}^5 E_2 + \overline{x}^2 E_5;
\end{aligned} \tag{27}$$

on the submanifold H.

Using (11), (16) and (20) we get

$$A_{E_3}X = -m\overline{x}^2 E_1 - m\overline{x}^1 E_2; \quad A_{E_4}X = -\frac{1}{2}\left(\frac{3}{2}a\overline{x}^2 + m\overline{x}^5\right)E_2 + \frac{1}{2}m\overline{x}^2 E_5;$$

$$\gamma(X) = \frac{\sqrt{3}}{2}\left(a\overline{x}^2 - m\overline{x}^5\right).$$

Then the formulas of Gauss and Weingarten (11) become

$$\overline{\nabla}_X Y = \nabla_X Y - m\left(\overline{x}^2\overline{y}^1 + \overline{x}^1\overline{y}^2\right)E_3 + \frac{1}{2}\left(\left(\frac{3}{2}a\overline{x}^2 + m\overline{x}^5\right)\overline{y}^2 + m\overline{x}^2\overline{y}^5\right)E_4;$$

$$\overline{\nabla}_X E_3 = m\overline{x}^2E_1 + m\overline{x}^1E_2 + \frac{\sqrt{3}}{2}\left(a\overline{x}^2 - m\overline{x}^5\right)E_4;$$

$$\overline{\nabla}_X E_4 = \frac{1}{2}\left(\frac{3}{2}a\overline{x}^2 + m\overline{x}^5\right)E_2 - \frac{1}{2}m\overline{x}^2E_5 + \frac{\sqrt{3}}{2}\left(a\overline{x}^2 - m\overline{x}^5\right)E_3.$$

Having in mind the last formulas, (26) and (27) we compute the tensor F of H

$$F(X,Y,Z) = -\frac{\sqrt{3}}{2}a\overline{x}^2(\overline{y}^1\overline{z}^2 + \overline{y}^2\overline{z}^1), \quad X,Y,Z \in \chi(H)$$

and verify that the submanifold (H,ϕ,ξ,η,g) of G, where (ϕ,ξ,η) is defined by (27) is an almost contact manifold with Norden metric in the class $\mathcal{F}_4 \oplus \mathcal{F}_8$.

References

1. D. Blair. *Contact manifolds in Riemannian geometry*, Lecture Notes in Math., vol. 509, Springer Verlag, Berlin, 1976.
2. F. Warner. *Foundations of differentiable manifolds and Lie groups*, Springer Verlag, New York Berlin Heidelberg Tokyo, 1983.
3. G. Ganchev, V. Mihova and K. Gribachev. *Almost contact manifolds with B-metric*, Math. Balkanica, vol. 7, 1993, 262–276.
4. G. Nakova, K. Gribachev. *Submanifolds of some almost contact manifolds with B-metric with codimension two, I*, Math. Balkanica, vol. 11, 1997, 255–267.
5. G. Nakova, K. Gribachev. *Submanifolds of some almost contact manifolds with B-metric with codimension two, II*, Math. Balkanica, vol. 12, 1998, 93–108.
6. G. Nakova. *On some non-integrable almost contact manifolds with Norden metric of dimension 5*, Topics in contemporary differential geometry, complex analysis and mathematical physics, Proceedings of the 8th International Workshop on Complex Structures and Vector Fields, Singapore, 2007, 252–260.